

Anisotropic Ordering in a Two-Temperature Lattice Gas

Attila Szolnoki and György Szabó

Research Institute for Materials Science, H-1525 Budapest, POB 49, Hungary

We consider a two-dimensional lattice gas model with repulsive nearest- and next-nearest-neighbor interactions that evolves in time according to anisotropic Kawasaki dynamics. The hopping of particles along the principal directions is governed by heat baths at different temperatures. The stationary states of this non-equilibrium model are studied using a generalized dynamical mean-field *medskip*analysis. In the stable state the particles form parallel chains which are oriented along the direction of the higher temperature. The resulting phase diagram is confirmed by Monte Carlo simulations.

PACS numbers: 05.50.+q, 05.70.Ln, 64.60.Cn

Systems driven into a non-equilibrium steady state can reveal a number of features not characteristic to the equilibrium state. An example is the well-known two-dimensional lattice gas model with attractive nearest-neighbor interaction whose degenerate ground states violate the $x - y$ symmetry of the system. Below a critical temperature the disordered (high temperature) phase segregates into a high- and a low-density phase and the strip region of the condensed particles can be oriented either horizontally or vertically if periodic boundary conditions are imposed. There are several different ways to provide non-equilibrium conditions and to violate the $x - y$ symmetry. In driven lattice gases introduced by Katz *et al.* [1] the particle jumps are under the influence of a uniform external electric field whose direction is chosen to be parallel to one of the principal axes [2]. Another way to define anisotropic dynamics is the so called two-temperature models are investigated extensively [3–6]. In these former models the particle jumps are coupled to two different heat baths, that is, the Kawasaki (exchange) dynamics is characterized by T_x and T_y temperatures along the x and y axes. In contrary to the previous dynamics, here there is no a uniform particle current through the system. At the same time, an energy transport exists from one of the heat baths to the other one driving the system into a non-equilibrium steady state. This approach can describe the effect of an alternating field (or polarized light) on the ordering process [2].

In the attractive interacting system the symmetry breaking is represented by the orientation of the interface separating the high- and a low-density phases. Significantly different symmetry breaking can be observed in the model discussed by Sadiq and Binder [7] which is characterized by repulsive nearest- and next-nearest-neighbor interactions with equal strength. In the half-filled system the particles form alternately occupied and empty columns (or rows). That is, here the symmetry breaking appears directly (as a bulk feature) in the four-fold degenerate ground states. The ordering process in this model is well understood [7–9] and the effect of the uniform driving field on the ordering process was also investigated [10]. It was found that the phases consisting of chains perpendicular to the external field become un-

stable and the stationary state is oriented along the field direction. In other words, the chain orientation of the ordered phase can be controlled by the application of an electric field with suitable direction. Similar effect is suspected when using an alternating field. This conjecture was the motivation of the present work. Now we will study the effect of anisotropic jump rate on the ordering process within the formalism of the two-temperature model.

First we introduce the model and investigate by adopting a simple dynamical mean-field theory [10]. Linear stability analysis has been performed to distinguish the stable and unstable solutions. In the light of these results the phase diagram is evaluated at higher level (4-point approximation) of dynamical mean-field theory. Finally the results are compared with Monte Carlo (MC) data.

We consider a half-filled, lattice-gas model with isotropic nearest- and next-nearest-neighbor repulsive interactions with equal strength on a square lattice. As usual the coupling constants as well as the Boltzmann constant are chosen to be unity. The kinetics is governed by Kawasaki dynamics characterized by single particle jumps to one of the empty nearest neighbor sites [11]. To avoid the difficulties arising from the non-analytic feature of Metropolis rate, Kawasaki hopping rate is used for directions $\alpha = x$ or y :

$$g_\alpha(\Delta H) = \frac{1}{1 + \exp(\Delta H/T_\alpha)} \quad , \quad (1)$$

where ΔH is the energy difference between the final and initial configurations. This jump rate is anisotropic and satisfies detailed balance at temperature T_y (T_x) in the vertical (horizontal) direction.

The equilibrium model ($T_x = T_y$) undergoes a continuous phase transition at a critical temperature $T_c = 0.525$. In the ground states the columns (or rows) are alternately occupied and empty providing fourfold degeneracy. Following the notation of Sadiq and Binder the lattice is divided into four interpenetrating sublattices and the 2×1 or 1×2 long-range order is characterized by the corresponding average sublattice occupations n_i ($i = 1, \dots, 4$) [7]. In the half-filled system the 2×1 states consisting of vertical chains are given as $n_1 = n_4 = (1 + m_x)/2$ and

$n_2 = n_3 = (1 - m_x)/2$, where $-1 \leq m_x \leq 1$ is the order parameter. Notice that m_x can be either positive or negative. Its absolute value as a function of temperature may be estimated by using the traditional mean-field approximation. Similar expressions are obtained for the 1×2 states of horizontal chains, namely $n_1 = n_2 = (1 + m_y)/2$ and $n_3 = n_4 = (1 - m_y)/2$. It is emphasized that the four possible states are equivalent ($|m_x| = |m_y|$) in the equilibrium system. Evidently, in the high temperature disordered phase $m_\alpha = 0$.

In the following simple dynamical mean-field approximation [10] we will suppose that the long-range order still exists under the present non-equilibrium condition. It means that the anisotropic jump rate results in only a difference between the solutions characterized by m_x and m_y .

According to Eq. (1) the time evolution of n_1 is given by summarizing the contributions of the jumps modifying the particle density in sublattice 1 as

$$\frac{dn_1}{dt} = -2n_1[(1 - n_2)g_x(E_{21}) + (1 - n_4)g_y(E_{41})] + 2(1 - n_1)[n_2g_x(E_{12}) + n_4g_y(E_{14})], \quad (2)$$

where E_{ij} is the average value of ΔH when a particle jumps from sublattice i to j . Similar equations can be derived for the rest of sublattice occupations. Assuming a 2×1 or 1×2 structures (vertical or horizontal chains) the stationary solution of Eq. (2) obeys the following general form:

$$\frac{1 + m_\alpha}{1 - m_\alpha} = \exp(3m_\alpha/2T_\alpha) \quad (3)$$

where $\alpha = x$ or y respectively. Notice that the jumps along the chains do not modify the particle densities within the chains therefore the value of the corresponding order parameter depends only on the perpendicular jump rate (or perpendicular temperature). The order parameter as a function of perpendicular temperature can be determined numerically. This simple approximation predicts continuous phase transition with a critical temperature $T_c^{(MF)} = 3/4$.

Obviously, we have two different solutions if $T_x \neq T_y$. In this situation the linear stability analysis is used to determine which solution remains stable. This method proved to be very useful when investigating the effect of the uniform driving field on these ordered structure in the same lattice-gas model [10]. Now we restrict ourselves to the survey of results without dealing with mathematical details given in a previous work [10]. It is found that the ordered structure appears when $\min(T_x, T_y) < T_c^{(MF)}$ and the chains in the stable phases are parallel to direction belonging to the higher temperature. As mentioned above, the value of the corresponding order parameter depends only on the perpendicular temperature. As a consequence the chain structure remains stable even in the limit case when the parallel temperature goes to infinity. The resultant phase diagram is illustrated in Fig. 1.




FIG. 1. Phase diagram predicted by the simplest mean-field theory. The arrows indicate the stable chain directions in the ordered regions.

The above results imply the possibility of reorientation in the low temperature region. Namely, exchanging the values of T_x and T_y a particle rearrangement is expected as it is observed by MC simulations under the effect of a uniform field [10]. It is rather difficult to realize the two-temperature model, but as we mentioned earlier an alternate electric field can cause similar behavior. Detailed analysis of the reorientation phenomenon goes beyond the scope of the present work. Instead, we focus our attention to have a more accurate phase diagram.

The prediction of the above simple mean-field theory can be improved by using the generalized mean-field theory at the levels of 2- and 4-point approximations. In these calculations we have to determine the probability of all the possible configurations on 2- and 4-point clusters by evaluating numerically the stationary solution of a set of equations which describe the time variation of configuration probabilities. Details of such a calculation are given in previous papers [12,13]. The 2-point (pair) approximation fails to reproduce the phase transition even in the equilibrium model ($T_y = T_x$). More precisely, this approximation predicts first-order transition. Similar discrepancies have already been observed for other systems [12,14]. This shortage of the present technique may be corrected by calculating with larger cluster(s).

Exploiting symmetries we have (only) four independent parameters to describe the probability of all the possible particle configurations on a square at the level of 4-point approximation. In equilibrium this method suggests a continuous order-disorder transition and the predicted critical temperature is $T_c^{(4p)} = 0.566$ which is close to the exact value mentioned above.

In the non-equilibrium case ($T_x \neq T_y$) the prediction of the 4-point approximation agrees qualitatively with those obtained above (see Fig. 1). However, now the critical temperature depends on both temperature. The numerical results are summarized in a revised phase diagram as plotted in Fig. 2. A weak anomaly has been

found when T_x and T_y are close to $T_c^{(4p)}$. Namely, the ordering temperatures *increase* first when moving away the equilibrium. After a low peak the transition temperature decreases toward a constant value, $T_c^{(4p)}(\infty) = 0.94T_c^{(4p)}$ if the parallel temperature goes to infinity. Hence, in agreement with the earlier approximation, the ordered state still exists even at infinite parallel temperature if the perpendicular temperature is low enough.




FIG. 2. Comparison of phase diagrams predicted by 4-point approximation (solid line) and MC simulations (squares).

To check the above predictions a series of Monte Carlo simulations [15] has been carried out varying the temperatures T_x and T_y . Using periodic boundary conditions the simulations are performed on a square system for different sizes $L = 20, 30, 40, 80$, and 200 . The size dependence of data has indicated no serious finite size effects. We have determined the time averages of the energy and order parameter varying the perpendicular temperature. These simulations have justified that the ordering process remains continuous for all the parallel temperatures we studied. From these data we could determine the value of critical temperatures. The results confirm the above theoretical predictions as demonstrated in Fig. 2. This figure visualizes that the transition temperature depends slightly on the parallel temperature, in a very similar way as suggested by the 4-point approximation and its value tends to $0.98T_c^{(MC)}$ when T_x or T_y goes to infinity.

In summary, we have studied the non-equilibrium ordering process in a two-dimensional lattice gas coupled to two thermal baths at different temperatures. In the equilibrium system the particles form parallel chains oriented either horizontally or vertically. Using a simple mean-field theory including linear stability analysis we have found that the ordered structure appears in the nonequilibrium case if the lower temperature is less than a critical value and the chains are oriented along the higher temperature. This picture has been confirmed by dynamical 4-point approximation and MC simulations. Despite the prediction of the simple mean-field theory the sophisticated methods have indicated that the transition tem-

perature (as well as the temperature dependence of order parameter) depends slightly on the parallel temperature.

The possibility of orientation (or reorientation) process may appear in other two-dimensional systems belonging to the family of Ashkin-Teller models [16]. The present two-temperature approach gives some insight into this phenomena even for higher dimensions.

The authors are grateful to Christof Scheele for careful reading of the manuscript. This research was supported by the Hungarian National Research Fund (OTKA) under Grant No. F-7240.

-
- [1] S. Katz, J. L. Lebowitz, and H. Spohn, *J. Stat. Phys.* **34**, 497 (1984).
 - [2] For a recent review, see B. Schmittmann and R. K. P. Zia, in *Phase Transition and Critical Phenomena* edited by C. Domb and J. L. Lebowitz (Academic, New York, 1995), Vol. 17.
 - [3] Z. Cheng, P. L. Garrido, J. L. Lebowitz, and J. L. Vallés, *Europhys. Lett.* **14**, 507 (1991).
 - [4] E. L. Praestgaard, H. Larsen, and R. K. P. Zia, *Europhys. Lett.* **25**, 447 (1994).
 - [5] K. E. Bassler and Z. Rácz, *Phys. Rev. Lett.* **73**, 1320 (1994); *Phys. Rev. E* **52**, R9 (1995).
 - [6] K. E. Bassler and R. K. P. Zia, *J. Stat. Phys.* **80**, 499 (1995).
 - [7] A. Sadiq and K. Binder, *Surf. Sci.* **128**, 350 (1983).
 - [8] A. Sadiq and K. Binder, *Phys. Rev. Lett.* **51**, 674 (1983); *J. Stat. Phys.* **35**, 517 (1984).
 - [9] H. C. Fogedby and, O. G. Mouritsen, *Phys. Rev. B* **37**, 5962 (1988); K. A. Fichtorn and W. H. Weinberg, *Phys. Rev. Lett.* **68**, 604 (1992).
 - [10] G. Szabó and A. Szolnoki, *Phys. Rev. A* **41** 2235 (1990); G. Szabó, A. Szolnoki, and G. Ódor, *Phys. Rev. B* **46** 11432 (1992).
 - [11] K. Kawasaki, in *Phase Transition and Critical Phenomena*, ed. C. Domb and M. S. Green (Academic, New York, 1972), Vol. 2.
 - [12] R. Dickman, *Phys. Rev. A* **38**, 2588 (1988); **41**, 2192 (1990).
 - [13] A. Szolnoki and G. Szabó, *Phys. Rev. E* **48**, 611 (1993).
 - [14] G. Ódor, N. Boccara, and G. Szabó, *Phys. Rev. E* **48**, 3168 (1993).
 - [15] For a review, see, O. G. Mouritsen, *Computer Studies of Phase Transition and Critical Phenomena* (Springer, 1984), and K. Binder and D. W. Hermann *Monte Carlo Simulation in Statistical Physics* (Springer, 1988).
 - [16] J. Ashkin and E. Teller, *Phys. Rev.* **64**, 178 (1943).